

# Analysis of Plant Remains from the 44PY449 site

prepared by:

Matthew E. Biwer  
and  
Amber M. VanDerwarker

Report Submitted to TCR Garrow Associates, Inc.

## Table of Contents

	page
Introduction	2
Recovery and Preservation Bias	2
Methods of Quantification	3
Laboratory Procedures	4
Basic Results	4
Conclusions	5
References Cited	7
Table 1 (Summary data for Late Woodland plants identified at 44PY449).	6
Table 2 (Ubiquity Values, Standardized Counts, and Relative Percents of Plant Taxa from 44PY449 plant assemblage)	6

Appendices provided in separate MS Excel files:

Appendix A (Flotation data by sample)

## Introduction

Archaeological plant and animal assemblages represent only a small fraction of what was originally used and deposited by humans in open-air settings. Natural and cultural taphonomic processes can significantly modify organic remains, resulting in recovered assemblages that differ dramatically from the original deposits. As archaeologists, we examine collections that have undergone a series of processes—from the original collection of plants and animals by humans from the natural environment, to food preparation, cooking, discard, animal and insect scavenging, burial, decay, and weathering, to the eventual recovery of food residues by archaeologists. As a result, these food remains cannot be interpreted at face value. Instead, standard methodological procedures for sampling, quantification, and analysis allows us to make sense of botanical assemblages. Here we report on the identification and analysis of the archaeobotanical assemblage from the 44PY449, a site in Virginia with occupations likely dating to the Early Woodland period (6 samples).

## Recovery and Preservation Bias

Formation processes, or the ways in which an artifact or feature entered the archaeological record, are an important consideration in archaeobotanical analysis. In general, seeds may enter the archaeological record: 1) through gathering, cooking, or other processing activities; 2) by being incidentally brought to the site alongside a comestible and thrown away; 3) having been gathered for non-food purposes and discarded as waste; 4) as an inclusion in dung used for building material or fuel; 5) or blown in or accidentally brought back as a rider (see Hubbard 1976). These scenarios provide cultural and environmental context for the entry of plants into the archaeological record. The mode in which a plant is collected, transported, processed, used, and discarded will influence its entry and abundance in the archaeological record (see Dennel 1976; Miksicek 1987; Minnis 1981).

We must also consider how the botanic remains were preserved, as preservation impacts the composition of the archaeobotanical record. Macrobotanical remains in exposed environments will decompose rapidly due to biological, chemical, and weathering processes. However, archaeological plant remains may be preserved at an archaeological site: 1) in an extremely dry environment; 2) in an extremely wet (anaerobic) environment; or 3) through carbonization (Pearsall 1988). Dry preservation (desiccation) occurs in areas where the continual absence of moisture inhibits the development and growth of bacteria, fungi, and other microorganisms that assist in decomposition. Recovering desiccated remains from archaeological excavations is rare, but when encountered they often provide a more complete inventory of plants than the carbonized assemblage. In wet environments, water saturation (waterlogging) will create conditions that inhibit the growth of microorganisms and thus slow decomposition, resulting in remarkably preserved botanical remains; in extremely dry or wet environments, the favorable conditions for preservation often create unique challenges, such as the recovery of a preponderance of dense materials from a site. Finally, preservation through carbonization, in which organic material is converted into an inorganic matter, is the most commonly encountered vector of plant preservation worldwide (e.g., Hastorf and Popper 1988; Wright 2010).

The type or portion of plant that is recovered is also relevant to a discussion of preservation and formation processes of botanic remains in archaeological sites. Comestibles and other food-related items that have non-edible parts (e.g., maize cobs, nutshell) are often discarded into the fire as fuel (see Minnis 1981) or used as a tool. The discarded inedible

portions may survive into the archaeological record depending on their density. For example, nutshell, which tends to be denser than the edible parts (e.g., nutshell, dense seeds), has a higher likelihood of surviving the process of carbonization and thus being preserved in the archaeological record than a maize cob because they are more fragile than nutshell. On the other hand, edible portions of plants that are often consumed whole (e.g., beans, maize kernels) or raw (e.g., fruits), are less likely to enter into the archaeological record, though accidents do occur and are patterned (see Yarnell 1982). In this situation, coprolites may provide the only direct evidence of a consumable/consumed taxon.

### Methods of Quantification

Quantitative methods in archaeobotany have developed significantly over the past several decades (see Hastorf and Popper 1988; VanDerwarker 2010). The most common methods used by archaeobotanists for recording and quantifying plant remains are raw (absolute) counts and weights. Absolute counts and weights are unstandardized data and may reflect differential preservation, sampling, local environmental conditions, or other factors. These measures are a useful way to display original data as it was collected by the archaeobotanist and may be used by other researchers for comparative analysis. However, raw counts and weights are not appropriate for direct evaluation due to problems of comparability between plant taxa because they do not control for preservation biases and sampling error (see Miller 1988; Popper 1988).

One way to avoid the problems of absolute counts and weights is by using the ubiquity measure (Pearsall 2000:212-16; Popper 1988:60-64). This method of standardization calculates the percentage of samples in which a taxon is present relative to the total number of samples. The taxon is considered present whether there are 1 or 1,000 specimens, and the same frequency score is given no matter the count. For example, if maize is present in 6 of 10 samples, it receives a ubiquity score of 60%. This is an excellent means for dealing with differential preservation, as plants that may be overrepresented or underrepresented due to taphonomic processes have the same influence when present. Ubiquity is also useful for investigating spatial and temporal patterns of plant use within similar contexts, though the results but may be less meaningful when comparing contexts of differential deposition or use.

Density is another useful standardizing measure that uses a constant variable, such as soil volume, to create a comparative ratio to assess the relative abundance of plants at the site (Miller 1988; Scarry 1986). To calculate density, the absolute count of plant taxa (numerator) is divided by the total soil volume collected from a sample, context, or site (denominator). Standardizing botanic data using density controls for differences in soil volume between samples and allows for the direct comparison of samples of unequal size. A basic assumption in using this measure is that the larger soil volume sampled, the greater likelihood that (rare) plant remains that will be recovered (Miller 1988:73).

Standardizing by soil volume, however, does not control for the range of non-plant related activities that contribute to the deposit from which the soil sample derives. In other words, the density measure does not consider plant remains in terms of plant-related activities, but rather in terms of all of the activities that are represented in the deposit. Thus, if the analyst is interested in determining the importance of a specific plant relative to the other plants in a sample or context, then density measures may be inadequate. Rather, standardizing by plant weight (standardized count) might be more appropriate (Scarry 1986). Unlike the density measure, standardizing by plant weight considers the contribution of a specific plant or category

of plants solely in terms of plant-related activities. As a result, a plant weight ratio more accurately reflects spatial and temporal differences in plant use. As a quantitative category, plant weight is a sum of weights recorded for all carbonized plant specimens per sample or context. Thus, for each sample, there is a total weight of plant material—this figure is the denominator used to standardize the variable of interest.

Overall, ratios are useful tools that overcome some of the problems of absolute counts. However, it is important to note that ratios reveal only the *relative importance* of plants within depositional contexts, not the contribution of resources to ancient diet (see Scarry 1986). For example, the recovery of 100 nutshells and 10 maize kernels does not necessarily represent evidence that nuts were more important than maize to the diet of residents of a given site; preservation and sampling biases prohibit paleoethnobotanists from making definitive determinations on whether certain taxa were more common or important than others. For the purposes of the present analysis, I focus on a basic qualitative summary of the data, using ubiquity, standardized counts, and relative percent contribution of the taxon to the total botanic assemblage simply because there are too few samples to perform any meaningful quantitative analysis.

### Laboratory Procedures

Both the light and heavy fractions of the flotation samples were analyzed. Although the materials from the light and heavy fractions were processed and sorted separately, data from the two fractions were combined for analysis. According to standard practice, the light fractions were weighed and then sifted through 2.0 mm, 1.4 mm, and 0.7 mm standard geological sieves. Carbonized plant remains from both fractions were sorted in entirety down to the 2.0 mm sieve size with the aid of a stereoscopic microscope (10–40 X). Residue less than 2.0 mm in size was scanned for seeds, which were removed and counted. In addition, taxa encountered in the 1.4 mm sieve that were not identified from the 2.0 mm sieve were also removed, counted, and weighed.

Botanical materials were identified with reference to the paleoethnobotanical comparative collection at the University of California, Santa Barbara (UCSB) paleoethnobotany lab, various seed identification manuals (Delorit 1970; Martin and Barkley 1961; Medsger 1966; Minnis (2003), which allowed us to identify the range of taxa native to the region. Taxonomic identification was not always possible—some plant specimens lacked diagnostic features altogether or were too highly fragmented. As a result, these specimens were classified as “unidentified.” Once the plant specimens were sorted and identified, counts, weights (in grams) and provenience information were recorded. Wood was weighed but not counted, and no wood identification was conducted. Other than counts and weights, no other measurements were taken on any specimens.

### Basic Results

This section presents the results of the identification of the carbonized plant remains from 44PY449. A total of six flotation samples from this site were sent to UCSB for analysis. Plant data from flotation samples are summarized in Table 1. Raw counts and weights are provided for each taxon; plant weight and wood weight are also provided. All samples were sorted, representing a total plant weight of 6.54 grams and a total wood weight of 6.18 grams. Combined, these samples yielded four plant taxa (all were identified to the Genus level), including nuts and seeds/leafy greens (Table 1). No crop taxa, such as maize (*Zea mays*), were

identified. Ubiquity, standardized count, and relative percent measures are provided in Table 2. Given the limited number of samples (n=6) from this site, no further quantitative analysis was conducted.

Nutshell recovered from the 44PY449 flotation samples includes hickory (*Carya* spp.) and walnut (*Juglans* sp.). Hickory was the most abundant nut recovered (n=50) with low amounts of walnut (n=2) also present. Hickory and walnut nuts require extensive processing before they are rendered palatable (Petrucci and Wickens 1984). The hickory nutmeat is so tightly enmeshed in the interior shell that picking the nutshells from the cracked shell casing is a time-consuming task. Instead, hickory nuts were generally pounded into pieces and boiled to extract the oil (Ulmer and Beck 1951). The process of boiling the pounded hickory nuts separates the pieces of shell, which sink to the bottom of the pot, from the oil, which rises to the top as the nutmeats dissolve and can be skimmed off or decanted. This oil/milk would then be used as an added ingredient in soups and stews, as a condiment for vegetables, or as a general sauce or beverage (see Scarry 2003; Talalay et al. 1984).

Two species were labeled as wild/miscellaneous seeds. These seeds, including purslane (*Portulaca* sp.) and carpetweed (*Mullugo* sp.), produce edible greens and/or seeds (Tables 1 & 2). In addition, several unidentifiable small carbonized fragments were also recovered.

### Conclusions

Overall, the flotation samples from 44PY449 produced low counts of plant remains, though this may be expected for such a small set of samples. These plants mainly represent food plants, greens, and/or herbs. The lack of cultivated species in the assemblage suggests that the site residents either did not practice horticulture or agriculture subsistence strategies at this locale, or the sampling strategies were inadequate in capturing the full range of deposited plant material. Given the data present, it is clear that some foraging activities occurred at the site, including the gathering of nuts, fruits, and wild greens, and possibly the management of local nut groves in the surrounding area.

Table 1. Counts and Weights of Plant Taxa by Sample

Site: VA-FS-13/44PY449	Sample Number	65		66		67		68		69		70	
	T.U.					5				5			
	Block	2		2		2		2		2		2	
	Feature	1		1		2		2		2		3	
	Half	S 1/2		N 1/2									
	Level							9		10			
	Plant Weight (g)	1		3.25		1.43		0.3		0.41		0.15	
	Wood Weight (g)	0.88		3.14		1.34		0.29		0.4		0.13	
Common Name	Taxonomic ID	Count	Grams	Count	Grams	Count	Grams	Count	Grams	Count	Grams	Count	Grams
<b>Nuts</b>													
Hickory	<i>Carya</i> sp.	12	0.11	9	0.08	25	0.21	3	0.01			1	0.01
Walnut	<i>Juglans</i> sp.					2	0.02						
<b>Wild/Misc.</b>													
Purslane	<i>Portulaca</i> sp.	1	0.01	1	0.01	6	0.02						
Carpetweed	<i>Mollugo</i> sp.			1	0.01								
<b>Unidentified</b>													
UID Seed				1	0.01	2	0.02						
Unidentified						4	0.04			1	0.01	3	0.01

Table 2. Ubiquity Values, Standardized Counts, and Relative Percents of Plants from 44PY449

Common Name	Taxonomic ID	Count	Grams	Ubiquity	Relative Percent	Standardized Count
<b>Nuts</b>						
Hickory	<i>Carya</i> sp.	50	0.42	83%	69.44	86.21
Walnut	<i>Juglans</i> sp.	2	0.02	17%	2.78	3.45
<b>Wild/Misc.</b>						
Purslane	<i>Portulaca</i> sp.	8	0.04	50%	11.11	13.79
Carpetweed	<i>Mollugo</i> sp.	1	0.01	17%	1.39	1.72
<b>Unidentified</b>						
UID Seed		3	0.03	33%	4.17	5.17
Unidentified		8	0.06	50%	11.11	13.79

## References Cited

Delorit, R. J.

1970 *Illustrated Taxonomy Manual of Weed Seeds*. Wisconsin State University, River Falls.

Dennell, R. W.

1976 The Economic Importance of Plant Resources Represented on Archaeological Sites. *Journal of Archaeological Science*, 3(3) pp:275–284.

Hastorf, Christine A., and Virginia S. Popper (editors)

1988 *Current Paleoethnobotany: Analytical Methods and Cultural Interpretations of Archaeological Plant Remains*. The University of Chicago Press, Chicago and London.

Hubbard, R.N.L.B

1976 On the strength of the evidence for prehistoric crop processing activities. *Journal of Archaeological Science*, 3(3) pp:257–265.

Martin, A. C., and W. D. Barkley

1961 *Seed Identification Manual*. University of California Press, Berkeley.

Medsgger, Oliver Perry

1966 *Edible Wild Plants*. Collier Books, New York.

Miksicek, Charles H.

1987 Formation Processes of the Archaeobotanical Record. In *Advances in Archaeological Method and Theory*, Vol. 10, edited by M. Schiffer, pp. 211–247. Academic Press, New York.

Miller, Naomi F.

1988 Ratios in Paleoethnobotanical Analysis. In *Current Paleoethnobotany: Analytical Methods and Cultural Interpretations of Archaeological Plant Remains*, ed. by C. H. Hastorf and V. S. Popper, pp. 72–85. The University of Chicago Press, Chicago and London.

Minnis, Paul E.

1981 Seeds in Archaeological Sites: Sources and Some Interpretive Problems. *American Antiquity* 46(1):143–152.

Minnis, Paul E. (editor)

2003 *People and Plants in Eastern North America*. Smithsonian Books, Washington and London.

Pearsall, Deborah

2000 *Paleoethnobotany: A Handbook of Procedures*. (2<sup>nd</sup> ed.) Academic Press, San Diego.

1988 Interpreting the Meaning of Macroremain Abundance: The Impact of Source and Context. In *Current Paleoethnobotany: Analytical Methods and Cultural Interpretations of Archaeological Plant Remains*, edited by C. H. Hastorf and V. S. Popper, pp. 97–118. The University of Chicago Press, Chicago and London.

Petruso, Karl M., and Jere M. Wickens

1984 The Acorn in Aboriginal Subsistence in eastern North America: A Report on Miscellaneous Experiments. In *Experiments and Observations on Aboriginal Wild Plant Food Utilization in Eastern North America*, edited by P. Munson, 360-378. Indiana Historical Society, Indiana.

Popper, Virginia S.

1988 Selecting Quantitative Measures in Paleoethnobotany. In *Current Paleoethnobotany: Analytical Methods and Cultural Interpretations of Archaeological Plant Remains*, edited by C. H. Hastorf and V. S. Popper, pp. 53–71. The University of Chicago Press, Chicago and London.

Scarry, C. Margaret

2003 Patterns of wild plant utilization in the prehistoric Eastern Woodlands. In *People and Plants in the ancient eastern North America*, edited by P. J. Minnis, pp. 50-104. Smithsonian Institution Press, Washington D.C.

1986 Change in Plant Procurement and Production during the Emergence of the Moundville Chiefdom. Unpublished Ph.D. dissertation, Department of Anthropology, University of Michigan, Ann Arbor.

Swanton, John R.

1944 *The Indians of the Southeastern United States*. Bureau of American Ethnology Bulletin 137. Government Printing Office, Washington.

Talalay, Laurie, Donald R. Keller, and Patrick J. Munson

1984 Hickory Nuts, Walnuts, Bitternuts, and Hazelnuts: Observations and Experiments Relevant to Their Aboriginal Exploitation in Eastern North America. In *Experiments and Observations on Aboriginal Wild Plant Utilization in Eastern North America*, edited by P. J. Munson, pp. 338-359. Indiana Historical Society, Indianapolis.

Ulmer, Mary and Samuel E. Beck

1951 *Cherokee Cooklore: Preparing Cherokee Foods*. Museum of the Cherokee Indian, Cherokee.



VanDerwarker, Amber M.

2010 Correspondence Analysis and Principal Components Analysis as Methods for Integrating Archaeological Plant and Animal Remains. In *Integrating Zooarchaeology and Paleoethnobotany: A Consideration of Issues, Methods, and Cases*, edited by Amber M. VanDerwarker and Tanya M. Peres, pp. 65–74. Springer Science+Business Media, New York.

Wright, Patti J.

2010 Methodological Issues in Paleoethnobotany: A Consideration of Issues, Methods, and Cases. In *Integrating Zooarchaeology and Paleoethnobotany: A Consideration of Issues, Methods, and Cases*, edited by Amber M. VanDerwarker and Tanya M. Peres, pp. 15-36.

Yarnell, Richard A.

1982 Problems of interpretation of archaeological plant remains of the eastern woodlands. *Southeastern Archaeology* 1(1):1–7.